

Relationships between reservoir water quality and catchment habitat type

Gough, Rachel; Cohen, Yvonne; Fenner, Nathalie; Cannon, Jonathan; Freeman, Christopher

International Journal of Environmental Technology and Management

DOI:

[10.1504/IJETM.2016.074799](https://doi.org/10.1504/IJETM.2016.074799)

Published: 01/01/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Gough, R., Cohen, Y., Fenner, N., Cannon, J., & Freeman, C. (2016). Relationships between reservoir water quality and catchment habitat type. *International Journal of Environmental Technology and Management*, 19(1), 16-39. <https://doi.org/10.1504/IJETM.2016.074799>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Relationships between reservoir water quality and catchment habitat type

Rachel Gough^{1*}, Yvonne Cohen¹, Nathalie Fenner¹, Jonathan Cannon², Christopher Freeman¹

¹ *School of Biological Sciences, Bangor University, Deiniol Road, Bangor, Gwynedd, LL57 2UW, UK*

² *Dŵr Cymru Welsh Water Head Office, Pentwyn Road, Nelson, Treharris, Mid Glamorgan, CF46 6LY, UK*

*Corresponding author: bss408@bangor.ac.uk.

Abstract

Numerous catchment characteristics including topography, geology, soil and vegetation are reported to exert a strong influence on mean surface water properties. The present study employs a geographical information system (GIS) approach to examine, for the first time, the relationship between reservoir water quality (dissolved organic carbon (DOC) concentration, colour, nitrate concentration and pH) and catchment Phase 1 Habitat coverage. Analysis was conducted on 2 occasions and at 2 different spatial scales. Numerous statistically significant correlations were identified, suggesting the use of Phase 1 Habitat data could help improve predictive models of surface water quality. The occurrence and strength of correlations varied seasonally in response, we argue, to temporal variations in hydrological regime and anthropogenic activity. The data also suggest that the proximity of habitat types to the reservoir is significant in affecting reservoir water quality. The findings are used to recommend suitable measures for drinking water companies to mitigate against water quality issues.

Key words: dissolved organic carbon; drinking water; catchment; Phase 1 Habitat; soil; geographical information system.

1. Introduction

The biogeochemical properties of surface waters are acquired, to a large extent, during the passage of water through the catchment due to the interaction of water with vegetation, soils and mineral layers. Various organic and inorganic compounds will be solubilised and transported downstream during runoff, influencing solute concentrations, pH and ionic strength (Stutter et al., 2006).

Although surface water quality exhibits temporal variations in response to weather events and seasonal drivers (Gergel et al., 1999; Scott et al., 1998; Soulsby et al., 2006), physical catchment characteristics including topography, geology, soil and vegetation type will, to a large extent, determine mean biogeochemical characteristics (Billett and Cresser, 1992; Clair et al., 1994; Holden et al., 2007; Hope et al., 1994; Sobek et al., 2007). Amongst these variables, soil type is widely considered to represent a dominant control on surface water composition and quality (Aitkenhead et al., 1999; Billett and Cresser, 1992; Hope et al., 1997; Soulsby et al., 2006; Stutter et al., 2006).

Though its development is strongly influenced by other catchment features including soil characteristics, habitat type may also be an important factor affecting surface water quality. Vegetation type influences catchment hydrology, primary production and organic matter inputs (Ordóñez et al., 2008; Zhang et al., 2011), which affect soil composition and chemistry and in turn, drainage water quality. Forested catchments for example, have been associated with the production of dissolved organic carbon (DOC)-rich drainage waters (Grayson et al., 2012; Hope et al., 1994) with differences in DOC concentration and flux also reported between different tree species (Chow et al., 2009; Fröberg et al., 2011; Gough et al., 2012). Wetland habitat coverage is also reported to be a strong predictor of surface water DOC concentration (Gergel et al., 1999; Hope et al., 1994).

Surface water characteristics can also be strongly affected by anthropogenic activity. For example, the application of agricultural fertilisers has been associated with significant leaching of nutrients (nitrates and phosphates) into surface waters (Badruzzaman et al., 2012). Elevated nutrient concentrations may in turn result in eutrophication and algal blooms (Correll, 1998; Freeman et al.,

2009; Hecky and Kilham, 1988; Vollenweider, 1968), which are particularly problematic in drinking water supplies (Smith, 1998). Liming of agricultural land has been associated with increased surface water pH (Hindar et al., 2003). Drainage of wetland habitats in an attempt to improve their economic value has also been linked to elevated colour and DOC concentrations in surface waters (Holden et al., 2004; Wallage et al., 2006).

The Phase 1 Habitat Survey of Wales, completed by the Countryside Council of Wales (CCW) in 1997, provides a record of habitat coverage and land use (Howe et al., 2005). In its digitised form, using geographical information system (GIS) software, the data offers a useful means of measuring the spatial extent of different habitat types within catchments. Since the classification scheme includes both natural habitats and anthropogenic features (e.g. arable land and improved grassland), the data holds significant potential for researchers concerned with investigating catchment influences on surface water quality. The characteristics of surface waters supplying drinking water treatment works (WTWs) is important for water companies which have a responsibility to provide a safe and reliable drinking water supply for their customers. The concentration of DOC in surface waters is particularly important, with the removal of DOC from drinking water supplies representing the single biggest treatment cost for the water treatment industry (Watts et al., 2001). Elevated DOC concentrations in raw water can inflate treatment costs by increasing the coagulant and disinfectant doses required (Chow et al., 2005; Edzwald, 1993) and the frequency of filter backwashes (Eikebrokk et al., 2004). DOC in finished water is problematic since it can cause undesirable colour, odour and taste (Davies et al., 2004; WHO, 2011), transports organic and inorganic micro-pollutants (Gao et al., 1998; Rothwell et al., 2007) and leads to bacterial regrowth in distribution systems (Prévost et al., 1998). Crucially, DOC also acts as a precursor to potentially harmful disinfection by-products (DBPs) including trihalomethanes (THMs). These are formed during chlorination, a treatment necessary to ensure that finished water meets microbiological safety standards (WHO, 2011).

Rising surface water DOC concentrations have been observed in many areas of central and northern Europe and North America in the past couple of decades (Freeman et al., 2001; Hejzlar et al., 2003; Monteith et al., 2007; Skjelkvåle et al., 2005; Stoddard et al., 2003; Worrall et al., 2003). In the UK, measurements undertaken at 22 upland sites showed a mean increase in DOC concentration of 91% between 1988 and 2003 (Evans et al., 2005). DOC concentrations tend to be highest, and rising most rapidly in peat-dominated, upland catchments (Freeman et al., 2001), which in the UK, supply over 70% of drinking water (Watts et al., 2001). In this context of declining surface water quality, developing a better understanding of catchment influences is crucial for the drinking water industry. The importance of catchment characteristics in affecting the quality of drinking water supplies is recognised by the UK drinking water regulator, the drinking water inspectorate (DWI) who recommend that “*catchment and raw water source protection*” is included in the drinking water safety plans of drinking water providers (DWI, 2005).

A GIS approach, which can offer an effective means of visualising and measuring landscape features is increasingly being used in the study of catchment influences on hydrochemistry. GIS software has become an important tool in the modelling of hydrological processes and its use in developing predictive models for various water quality parameters within catchments based on land use and other catchment characteristics is particularly relevant for water treatment companies. For example Foster and McDonald (2000) used GIS and spatially referenced data on pastoral farming intensity to model and display sources of cryptosporidium risk in drinking water catchments. Lake et al. (2003) developed a nitrate leaching model using GIS and information on a number of physical catchment characteristics. This was used to identify areas of groundwater vulnerable to nitrate pollution. Recently, Grayson et al. (2012) used a GIS approach, and ITE land cover data (similar to Phase 1 Habitat data) to identify correlations between drinking water reservoir colour and the spatial extent of different land cover classes. A multicriteria evaluation approach was then used to develop a predictive model for water colour production potential in the catchments and create a colorimetric,

risk-based map from the data. However, as yet, the use of Phase 1 Habitat data for predicting catchment water quality has not been explored.

This study investigates potential relationships between Phase 1 Habitat classes and reservoir water quality (DOC concentration, colour, nitrate concentration and pH). GIS mapping was used to measure the spatial extent of Phase 1 Habitat types in 16 drinking water reservoir catchments in north Wales. Correlation analysis was then used to identify statistically significant relationships between these land cover classes and reservoir water quality in spring and autumn. Analyses were carried out both at a whole-catchment scale, and in a 250 m buffer zone surrounding the reservoirs in order to assess the importance of proximity in the occurrence and strength of the correlations. Such research is important for informing future catchment management practices. Identifying problematic land cover will also help water treatment companies target monitoring programmes and mitigation strategies, and improved understanding of seasonality in raw water quality will also enable better optimization of treatment processes.

2. Methods

2.1. Study sites and sampling regime

Water samples were collected on 2 occasions (in September 2007 and March 2008) from the raw water (i.e. pre-treatment) supply of 16 WTWs in north Wales. Where the raw water supply was derived from more than 1 reservoir, composite samples were collected, and Phase 1 Habitat data was also combined. The timing of sampling was chosen to correspond with the seasonal maximum (autumn) and minimum (spring) in reservoir DOC concentration. 14 of the WTWs included in this study are located in upland catchments, with the remaining 2 situated in lowland, agricultural areas. Uplands are defined as areas more than 250 m above sea level (Mitchell, 1991). These areas are typically characterised by high rainfall, low mean temperatures and acidic soils (Foster and McDonald, 2000).

2.2. Hydrochemical analysis

pH was measured on un-filtered samples using a Mettler Toledo S20 pH meter (Mettler Toledo, Leicester, UK), calibrated daily with pH 4 and pH 7 reference standards (Sigma-Aldrich, Dorset, UK).

Colour measurements (Hazen) were obtained from WTW data at the time of sample collection. One degree Hazen ($1 \text{ mg L}^{-1} \text{ Pt/Co}$) is defined as the colour produced by $1 \text{ mg L}^{-1} \text{ Pt}$ (as K_2PtCl_6) in the presence of 2 mg L^{-1} cobalt (II) chloride hexahydrate (Mitchell and McDonald, 1992).

Before DOC measurement, samples were passed through a $0.45 \mu\text{m}$ cellulose acetate filter to remove particulate organic carbon, as per the operational definition of DOC (Thurman, 1985). DOC concentrations were determined using a Shimadzu Total Organic Carbon 5000 analyser (Shimadzu, Milton Keynes, UK), with a carrier gas of high purity air at a flow rate of 150 mL min^{-1} and a $33 \mu\text{L}$ injection volume. Calibration was performed with a one point calibration, using a $100 \text{ mg L}^{-1} \text{ KO}_4\text{H}_5\text{C}_8$ solution (total organic carbon – TOC) and a $100 \text{ mg L}^{-1} \text{ Na}_2\text{CO}_3/100 \text{ mg L}^{-1} \text{ NaHCO}_3$ solution (inorganic carbon – IC). DOC concentrations were calculated by subtracting IC values from TOC values. Analysis of TOC and IC standard solutions at 10 mg L^{-1} intervals demonstrated that the analyser performed linearly from 0 to 200 mg L^{-1} , with r^2 values > 0.9 . All reagents were supplied by Sigma-Aldrich, Dorset, UK.

Nitrate concentration was determined using a Dionex DX-120 ion chromatograph equipped with an IonPac AS14A anion analytical column (both Thermo Scientific, Hertfordshire, UK). The eluent was a $1.0 \text{ mM Na}_2\text{HCO}_3/8.0 \text{ mM NaCO}_3$ solution (reagents supplied by Sigma-Aldrich, Dorset, UK) made with Milli Q water and the flow rate, 1 mL min^{-1} . Concentrations were determined using a five point calibration with standard Dionex solutions.

2.3. Geographical information systems (GIS) analysis

Version 9.2 of the ArcGIS package (ESRI, Buckinghamshire, UK) was used to display and quantify the spatial extent of habitat types within each reservoir catchment. First, the watersheds associated

with each reservoir were mapped. This was achieved using the *Hydrology* functions in the *Spatial Analyst* extension and a digital elevation model downloaded from Digimap (EDINA, 2014) (10 m resolution). Defined watersheds were then clipped to other GIS layers displaying habitat type. Habitat information was displayed using digitised version of the Phase 1 Habitat Survey of Wales (Howe et al., 2005). In addition to this whole-catchment analysis, habitat coverage was also measured in a 250 m-wide zone around the perimeter of each reservoir.

2.4. Statistical analysis

For statistical analysis, Phase 1 Habitat categories were organized into more generalised groupings (Table 1). Statistical analysis was performed using version 20 of the SPSS statistical package (IBM, New York, USA). Depending on the conditions satisfied by the data, Pearson's correlation and Spearman's correlation analyses was employed to test for significant correlations between Phase 1 habitat type coverage and reservoir water quality. This analysis was also performed using the subset of Phase 1 Habitat data covering a zone of 250 m directly adjacent to the reservoir.

3. Results and discussion

3.1. DOC and colour

The absence of any statistically significant correlations between catchment woodland and scrub coverage and reservoir DOC concentration and colour (Table 2 and 3) is surprising given that previous research indicates a strong positive relationship between forest coverage and DOC concentration (Grayson et al., 2012; Hope et al., 1994). High DOC flux from forested catchments is partly due to high DOC loading as rainwater passes through above ground biomass (Kawasaki et al., 2005; Stevens et al., 1989) as well as the large source of leachable carbon in the litter layer (Hongve, 1999). However, DOC concentrations are also reported to vary significantly between different tree species (Gough et al., 2012). Our habitat categories did not account for this potential variation, which may explain the absence of any statistically significant correlations in this study.

A moderate negative correlation was observed between unimproved grassland and spring DOC concentration at the whole catchment scale ($p < 0.05$; Table 2) and no correlations between unimproved grassland and DOC or colour in the 250 m buffer zone analysis (Table 3). The negative correlation corroborates the findings of previous studies. For example, Grayson et al. (2012) report a significant negative correlation between water colour and moorland grass coverage across 18 drinking water catchments in Yorkshire. In a UK-wide study, Armstrong et al. (2007) found that heather dominated, drained catchments produced the highest water colour followed by mixed vegetation and grass dominated catchments. Van den Berg et al. (2012) also reported lowest mean pore water DOC concentration in grassland sites compared with other vegetation categories (woodlands, heathlands and moorlands) in their survey of 41 UK sites. This association may relate to solubility controls since colour release in temperate grasslands is reported to be suppressed by acidic conditions (Hopkins et al., 1990; Miller, 2008). However, our unimproved grassland category included neutral and calcareous grassland and no correlations were identified in the present study between pH and unimproved grassland coverage.

Negative correlations were identified between tall herb and fern habitat coverage and autumn DOC concentration and colour at the whole catchment scale (both $p < 0.05$; Table 2) and between tall herb and fern coverage and autumn colour in the 250 m buffer zone analysis ($p < 0.05$; Table 3). At first this result seems unexpected since bracken coverage, which was dominant in this habitat class, has been associated with high primary productivity and the accumulation of large amounts of litter, forming a large pool of organic matter (Marrs et al., 2000). However, Potthast et al. (2012) observed that, compared with pasture land (*Setaria* grass) the litter present in bracken habitat showed a significantly lower rate of decay. In addition, they found a significant decrease in microbial biomass and activity when pasture land was invaded by bracken. Therefore, if bracken coverage tends to replace improved grassland habitats in drinking water catchments, its presence may reduce DOC production. The occurrence of these correlations in autumn may relate to this being the litter fall

period, when the leaching of DOC from decomposing litter would normally contribute significantly to DOC export (Kalbitz et al., 2000).

Negative correlations between heathland coverage and DOC concentration and colour occurred at both spatial scales. In the whole catchment analysis, heathland coverage displayed a moderate negative correlation with autumn DOC concentration and colour (both $p < 0.05$) and a strong negative correlation with spring colour ($p < 0.01$; Table 2). In the 250 m buffer analysis a moderate negative correlation was identified with autumn DOC concentration and colour (both $p < 0.05$), a moderate negative correlation with spring DOC concentration ($p < 0.05$) and a strong negative correlation with spring colour ($p < 0.01$; Table 3). These negative relationships were surprising given that *Calluna*, a common species in heath habitats, has been reported to produce highly-coloured drainage water (Grayson et al., 2012). This has been attributed to their relatively dry soil conditions which confer high rates of aerobic microbial decomposition (Clutterbuck and Yallop, 2010). Many heath habitats were also formed as a result of peatland drainage. This former status is likely to further enhance DOC and colour release due to the large carbon stocks associated with peat substrate (Fenner et al., 2009). Conversely however, moisture constraints in heath habitats are reported to inhibit phenol oxidase activity (Toberman et al., 2008). According to the enzymic latch theory, this can suppress DOC production by causing an accumulation of phenolic compounds which inhibit the activity of hydrolase enzymes (Freeman et al., 2001). This may explain the negative correlations between heathland coverage and reservoir DOC concentration and colour observed in the present study. Overall, the relationship appeared stronger at the 250 m buffer scale, suggesting that proximity to the reservoir affected the degree to which this habitat influenced reservoir water quality.

At the whole catchment scale a moderate positive correlation was identified between fen/ mire habitat and autumn DOC concentration ($p < 0.05$; Table 2). This habitat also correlated positively with autumn and spring DOC concentration in the 250 m buffer analysis (both $p < 0.05$; Table 3).

Positive correlations between swamp coverage and reservoir DOC concentration and colour were also identified and were striking in terms of the strength of the correlations observed and their occurrence at both spatial scales and both sampling times (Table 2 and 3); all were strong positive correlations ($p < 0.01$) except for the swamp/ autumn DOC concentration correlation at the whole catchment scale which was a moderate positive trend ($p < 0.05$). These positive relationships are likely to be linked to the wetland status of these habitats. Percentage wetland coverage has been identified as an important predictor of stream water DOC concentration (Eckhardt and Moore, 1990; Gergel et al., 1999; Hope et al., 1994). A combination of high primary productivity and low decomposition rates causes the accumulation of deep layers of peat in wetland environments (Mitsch and Gosselink, 2000). The considerable depth of organic material in such environments provides a large pool of available carbon (Thurman, 1985) and the inhibitory effect of anaerobic conditions on microbial metabolism promotes the formation of DOC end products (Fenner et al., 2009). In addition, in wetland systems, the depth of the organic horizon limits contact between drainage waters and the adsorption sites within the mineral soil horizon, which also contributes to high DOC loading (Tipping et al., 1999). However, our data also show an absence of statistically significant correlations between DOC concentration/ colour and other habitat categories which are, or include, wetlands (marsh/ marshy grassland, bog, flush and spring). This suggests that the type of wetland present may be an important determinant of drainage water DOC concentration. It may be significant that, of these wetland habitat types, swamp and fen/ mire habitats tend to be more nutrient-rich than the other wetland habitats (Mitsch and Gosselink, 2000) which may support higher rates of primary productivity and thus a larger pool of organic carbon.

Strong positive correlations between arable coverage and DOC concentration were observed at the whole catchment scale in autumn and spring (both $p < 0.01$; Table 2). Moderate positive correlations were also identified with colour at both sampling times (both $p < 0.05$). The 250 m buffer zone could not be included in this analysis since there was only 1 catchment where arable land was present in this zone (Figure 1). The interpretation of these positive correlations is not straightforward since in

previous studies arable land use has been associated with lower carbon content than other land use types. For example, soil solution carbon concentrations for soils in northern Saskatchewan, Canada, decreased in the following order: aspen forest > recently cleared forest > wheat/ fallow field (McFee and Kelly, 1995). Similarly, in their review article, Chantigny (2003) reports that dissolved organic matter concentrations vary as follows: forest soils > grassland soils > arable soils. This variation, it is suggested, is partly due to differences in vegetation type (e.g. tree vs. herbaceous plant) (Chantigny, 2003) as well as the lower carbon content associated with arable soils (Zsolnay, 1996). In addition, aerobic conditions, which tend to occur in arable soils encourage the complete mineralisation of organic matter to CO₂, as opposed to DOC and CO₂ end products in anaerobic decomposition (Boddy et al., 2008; Fenner et al., 2009). However, water soluble carbon content in arable soils is also reported to vary depending on crop plants used (Zsolnay, 1996) and temporally, during crop cycles (Campbell et al., 1999) and with successive cultivations (Delprat et al., 1997). In addition, application of organic fertilisers on agricultural soils is reported to substantially increase the concentration of soluble organic carbon (Gregorich et al., 1998). Although it is not possible to isolate the cause of the positive correlations observed here between arable land use and DOC concentration/ colour, it is notable that the correlations occurred despite this land use being virtually absent in the reservoir 250 m buffer zone.

At the whole catchment scale moderate positive correlations were identified between buildings coverage and autumn DOC concentration ($p < 0.05$) and between the “other” category and autumn DOC concentration ($p < 0.05$) and spring colour ($p < 0.05$; Table 2). A moderate positive correlation was also found between buildings coverage and autumn DOC concentration in the 250 m buffer zone analysis ($p < 0.05$; Table 3). Given the rural location of the catchments in the present study it is likely that farm buildings will account for a significant proportion of the buildings category. Indeed, a strong positive correlation between buildings and arable coverage was identified in the whole catchment data ($r_s = 0.803$, $p < 0.01$). This correlation may therefore explain the relationship between buildings and DOC concentration, though the reason for this being confined to the autumn

analysis is unclear. The correlations between the “other” category and DOC concentration and colour at the whole catchment scale are also difficult to interpret since this category includes unknown habitat classes (“not accessed” land and “illegible” data inputs). It may be significant however, that bare ground (J.4; Table 1) is included in this category, which may provide a source of readily-leachable organic matter.

3.2. Nitrate and pH

Strong positive correlations were observed between arable coverage and nitrate concentration in the whole catchment analysis in autumn and spring (both $p < 0.01$; Table 2). This is likely to be caused by the leaching of organic or inorganic fertiliser (Neill, 1989). The stronger correlation in the spring analysis may be due to the timing of fertiliser application, which for arable crops tends to occur in late winter/spring (MAFF, 2000; Trudgill et al., 1991). As mentioned earlier, the 250 m buffer zone could not be included in the analysis due to there being only 1 catchment where arable was present in this zone. It is interesting therefore that strong correlations exist at the whole catchment scale despite arable coverage being virtually absent in the 250 m buffer zone. The application of fertiliser may also explain the strong positive correlations between improved grassland coverage and nitrate concentration in both the whole catchment analysis ($p < 0.01$ in autumn and spring; Table 2), and the 250 m buffer zone analysis ($p < 0.05$ and $p < 0.01$ in autumn and spring, respectively; Table 3). Again, stronger correlations in the spring analysis at both spatial scales are likely to be due to the timing of fertiliser application. The strong positive correlations between woodland and scrub coverage and spring nitrate concentrations at both spatial scales (both $p < 0.01$; Table 2 and 3) may be explained by the application of fertiliser prior to tree planting in commercial forestry plantations (Drinan et al., 2013).

Moderate positive correlations were identified between fen/ mire coverage and reservoir nitrate concentration in spring sampling at the whole catchment scale ($p < 0.05$; Table 2) and in both autumn and spring in the 250 m buffer analysis (both $p < 0.05$; Table 3). These correlations are likely

297 to relate to the nutrient status of this habitat; fen systems are typically associated with relatively
298 high nutrient concentrations due to their being supplied by drainage water from surrounding
299 mineral soil (Mitsch and Gosselink, 2000).

300 The positive correlation between buildings and nitrate concentration at the whole catchment scale
301 in spring ($p < 0.05$; Table 2) may be due to the positive correlation mentioned earlier between
302 buildings and arable coverage. In addition, the urine and droppings of mammals and birds has been
303 identified as an important non-agricultural source of ammonia (DEFRA, 2002). Nitrifying bacteria in
304 the soil may then convert ammonia to nitrate. Therefore, assuming that a significant proportion of
305 the buildings in this category are farms, then the leaching of ammonia from domestic animals may
306 also account for this correlation. The leaching of nitrates from septic tanks and fertiliser stores may
307 also explain this association.

308 A moderate positive correlation was observed between marsh/ marshy grassland coverage and
309 reservoir nitrate concentration but only in the 250 m buffer analysis in spring ($p < 0.05$; Table 3). The
310 reason for this is not clear but may be an artefact of the positive association between marsh/
311 marshy grassland and other habitat types displaying a positive correlation with nitrate. For example,
312 in the 250 m buffer zone analysis, marsh/ marshy grassland coverage correlates positively with
313 woodland and scrub ($r_s = 0.646$, $p < 0.01$), improved grassland ($r_s = 0.771$, $p < 0.01$) and fen/ mire
314 habitat ($r_s = 0.560$, $p < 0.05$), all of which show a positive correlation with spring nitrate
315 concentration at this spatial scale.

316 The supply of nitrate and phosphate is critical in determining the growth rates of phytoplankton in
317 freshwater systems with elevated concentrations resulting in eutrophication in some cases (Hecky
318 and Kilham, 1988). In drinking water sources algal blooms can lead to a number of treatment issues
319 including taste and odour problems, elevated TOC levels, increased coagulant and chlorine demand,
320 membrane fouling and an increase in DBPs (Bernhardt et al., 1991; Li et al., 2012; Nguyen et al.,
321 2005). Elevated reservoir nitrate concentrations may also increase the formation of nitrogenous

DBPs (NDBPs), produced during the disinfection stage of water treatment either directly, or indirectly *via* increased algal biomass and consequently increased concentrations of dissolved organic nitrogen in the raw water (Ritson et al., 2014).

The negative correlation between heathland coverage and reservoir pH in the 250 m buffer analysis ($p < 0.05$; Table 3) is likely to be related to the preference of heath vegetation for acidic soils (Holden et al., 2007) which has a corresponding effect on drainage water pH (Cresser and Edwards, 1987). A positive relationship has been reported between DOC solubility and pH (Lumsdon et al., 2005). Thus solubility controls may also help to explain the negative correlations between heathland coverage and reservoir DOC concentration and colour.

The absence of correlations between pH and some of the other habitat types which tend to be associated with peat substrates (bare peat, bog, fen/ mire, flush and spring and swamp) is surprising given that peatlands tend to produce acidic drainage waters. This is reported to result from the accumulation of organic acids, the enhanced activity of sulphur-metabolising bacteria under waterlogged conditions and high cation exchange capacity (Clymo, 1964; Urban et al., 1995).

Coniferous forest stands, which represent a large proportion of forest coverage in north Wales, are also associated with acidic drainage waters (Eisalou et al., 2013; Gough et al., 2012). A significant decrease in pH has been reported as rainwater passes through coniferous canopies and litter (Eisalou et al., 2013), due to the high exchangeable acidity of coniferous foliage and litter and the fact that coniferous litter is readily leached of organic acids (Alfredsson et al., 1998; Nykvist, 1963).

3.3. Temporal and spatial variations in correlations

At the whole catchment scale, more associations between Phase 1 Habitat categories and DOC concentration were identified in autumn than in spring. A difference in hydrological regime due to higher rainfall in September than March may explain this contrast. Higher rainfall will result in a larger contribution of surface runoff to discharge water (Horton, 1933) which is likely to enhance the

influence of surface characteristics such as vegetation/ litter characteristics. The influence of habitat may also be enhanced by higher above ground biomass following the growing season.

Overall there were fewer statistically significant correlations identified between habitat types and reservoir nitrate concentration in autumn compared with spring (Table 2 and 3). We have already suggested that fertiliser application may be significant in explaining a number of correlations between habitat type and reservoir nitrate concentration (Neill, 1989), and that its timing may explain the greater number and strength of correlations in spring (MAFF, 2000; Trudgill et al., 1991). However, the drivers of seasonal variations in surface water nitrate concentration are known to be complex, comprising numerous biogeochemical and hydrological processes (Martin et al., 2004). Stream nitrate concentrations tend to exhibit a summer minima and a winter maxima (Neill, 1989; Reynolds et al., 1992). This is explained in part by variations in the supply of nitrogen. For example, in the summer, the availability of leachable nitrate in the soil is limited by lower atmospheric inputs and plant uptake and in streams by macrophyte uptake (Cooke and Cooper, 1988) and denitrification (Hill, 1979). In winter on the other hand, plant uptake decreases, atmospheric inputs increase and in-stream losses decrease due to lower primary productivity (Reynolds et al., 1992). Surface water nitrate levels may also be transport-limited, with a strong association reported with precipitation and discharge (Neill, 1989; Trudgill et al., 1991). However, given that lower rainfall totals were recorded in March than in September, it is more likely that reservoir nitrate levels in spring (which were higher than in autumn), were supply-limited.

It is difficult to interpret the overall effect of spatial scale on relationships between habitat classes and reservoir water quality since at the 250 m buffer scale, a number of habitat classes are present in only a few catchments, or are absent altogether. For example, there was only 1 catchment where arable land was identified in the 250 m buffer zone. However, there was an obvious similarity in the occurrence of significant correlations at the 2 spatial scales. Although there was no clear difference in the strength of the correlations between the 2 spatial scales, this similarity would suggest that

371 Phase 1 Habitat coverage in the 250 m buffer zone was more important than in the wider catchment
372 in affecting reservoir water quality. Previous studies have reported improved regressions between
373 land use and surface water quality parameters when the riparian area was included, or weighted
374 more heavily than other areas (Levine and Jones, 1990; Osborne and Wiley, 1988).

375 The relationship between reservoir water quality and catchment characteristics at different spatial
376 scales may also vary temporally. For example, Gergel et al. (1999), in their study of wetland influence
377 on DOC concentrations in Wisconsin lakes and rivers, found that in autumn, wetland coverage in the
378 whole catchment was the best predictor of lake water DOC concentration whereas in summer,
379 wetland coverage within 50 m of lakes was the best predictor. This could relate to seasonal
380 hydrological changes since the timing of sampling in autumn and spring corresponded with base
381 flow and peak flow conditions, respectively (Hurley et al., 1995).

382 In this study we noted the absence of a number correlations between Phase 1 Habitat classes and
383 surface water parameters that would typically be expected. For example the lack of positive
384 correlations between woodland and scrub habitat and a number of wetland habitats and DOC
385 concentration/ colour was unexpected. This may be due to the influence of various other catchment
386 features not included in the present study, but which previous studies have reported to influence
387 surface water chemistry. For example, slope will mediate the relationship between catchment
388 characteristics and surface water quality due to its influence on surface runoff (Rochelle et al., 1989)
389 as well as being a predictor of soil organic horizon depth (Rasmussen et al., 1989) and wetland
390 abundance (Eckhardt and Moore, 1990). The development of a particular soil type reflects a number
391 of factors including climate, parent material, topography and vegetation and is reported to be a
392 crucial factor in determining surface water composition and quality (Aitkenhead et al., 1999; Billett
393 and Cresser, 1992; Hope et al., 1997). Indeed information on soil chemical characteristics has formed
394 the basis of a number of predictive models for stream water solute concentrations (Billett and
395 Cresser, 1992; Christophersen and Wright, 1981; Cosby et al., 1985). Soil type influences spatial

patterns of water flow and storage (Grayson and Western, 2001; Weiler and Naef, 2003). In addition, adsorption processes in mineral soils regulate the transport of organic carbon and the soil organic pool is reported to be the main factor controlling DOC flux in streams (Aitkenhead et al., 1999). It should also be noted that the topographic watershed does not necessarily correspond with groundwater influence, which may also strongly impact on reservoir water quality (Garrison et al., 1987). In addition, the Phase 1 Habitat Survey of Wales was conducted between 1987 and 1997 and it is likely that a number of land use changes have occurred in this time (Stevens et al., 2004). Nonetheless, the present study has demonstrated that the data continues to be relevant to the study of surface water quality and the extent of its coverage represents a significant benefit for drinking water companies.

Various processes occurring in the water body will also affect surface water chemistry. For example, as mentioned earlier, seasonal variations in the uptake of nitrate in surface waters influence nitrate concentrations (Cooke and Cooper, 1988). DOC loss from reservoirs is reported to occur as a result of sedimentation and mineralisation processes (Algesten et al., 2004), with precipitation also affecting DOC concentrations *via* a dilution effect (Engstrom, 1987). Conversely, DOC may be produced within the water body (autochthonous DOC), potentially suppressing the relationship between DOC concentration and colour and terrestrial drivers. Nonetheless, a substantial number of correlations were identified between Phase 1 Habitat data and reservoir water characteristics in the present study. This, we suggest, relates both to the direct influence of vegetation/ land cover on runoff and drainage water quality and also to habitat classes being predictors of other physical characteristics such as peat soils or certain management practices.

3.4. Implications for potable water treatment

Though previous research has cited the relationship between catchment wetland coverage and surface water DOC and colour loading (Eckhardt and Moore, 1990; Gergel et al., 1999; Hope et al., 1994), our data suggest that wetland type may significantly affect the magnitude of this relationship.

The identification of positive associations between swamp and, to a lesser extent, fen/ mire habitats and DOC concentration/ colour, possibly the result of their nutrient supply (Mitsch and Gosselink, 2000), may justify monitoring the quality of drainage waters in these areas. Given that these habitat types also occupy very small proportions of the catchments included in this study (Figure 2), it may be that diverting drainage water from these areas would be a cost-effective strategy for improving reservoir water quality. Monitoring of drainage waters and diversion of water courses may also be appropriate for areas of arable land use which arguably exerted the strongest influence on surface water quality. This land use class correlated with DOC, colour and nitrate concentrations at both sampling times despite being virtually absent from the 250 m buffer zone of the reservoirs. The apparent impact of arable land on reservoir water quality highlights the importance of excluding this activity from areas close to the reservoir. In cases where the diversion of problematic drainage waters is not possible, it may be appropriate to blend reservoir water with water from another catchment, as has been employed previously as a strategy to reduce water discolouration from peat (Grayson et al., 2012).

Allowing the expansion of habitat types whose coverage correlates negatively with DOC/ colour may be a suitable strategy in some cases. However, the potential benefits to surface water quality may be outweighed by other detrimental impacts. For example, tall herb and fern coverage, which in the present study was dominated by bracken, correlates negatively with DOC and colour but bracken habitat has no economic value and is associated with the leaching of carcinogenic compounds such as ptaquiloside (Rasmussen et al., 2003). Heathland coverage also correlated negatively with DOC concentration and colour in the present study, but this we argue, may relate to site-specific factors such as soil moisture constraints since *Calluna* vegetation is typically associated with highly-coloured waters (Grayson et al., 2012).

Given the number of statistically significant correlations identified in the present study, and the national scale of Phase 1 Habitat data, we suggest that future research should explore integrating

Phase 1 Habitat data into predictive models for reservoir water quality. The present study has also highlights the fact that correlations between catchment characteristics and surface water quality may vary on a seasonal basis; an important consideration as researchers seek to develop more sophisticated predictive models.

4. Conclusions

This study has considered, for the first time, the use of catchment Phase 1 Habitat data for predicting reservoir water quality. Our analysis was conducted at two different spatial and temporal scales, to investigate the effect of season and the proximity of habitat types to the reservoir in affecting potential associations between habitat type and water quality parameters.

Numerous statistically significant correlations were observed between Phase 1 Habitat classes and reservoir water quality. These could be explained either by the direct impact of vegetation on drainage water or its association with other physical catchment characteristics or land management practices. Arable land cover appeared to have the most substantial impact on reservoir water quality, correlating strongly with DOC concentration, colour and nitrate concentration at both sampling times. This was despite arable land being virtually absent from the 250 m buffer zone.

The degree to which habitat classes affected reservoir water quality appeared to vary on a seasonal basis, with more correlations between habitat classes and DOC concentration in autumn, and between habitat classes and nitrate concentration in spring. However, a striking similarity was observed between correlations at the whole catchment scale and within the 250 m buffer zone. We therefore suggest that in general, the influence of habitat coverage on reservoir water quality parameters increases with proximity to the reservoir.

Although previous research has identified a link between wetland abundance and surface water DOC/ colour loading, our findings suggest that the type of wetland habitat present is also important. We found that swamp and fen/ mire habitats were the only wetland types which correlated with

reservoir DOC or colour. This specificity, we suggest, may relate to the high nutrient levels in these habitats which may support higher rates of primary production than other wetland types.

Based on the number and strength of correlations observed, we suggest that predictive models for surface water characteristics based on catchment characteristics could be improved by incorporating Phase 1 Habitat data. The findings of this study are important for drinking water companies concerned with maintaining finished water quality and may be of use in targeting monitoring of drainage water in catchments and selecting appropriate mitigation strategies such as diverting or blending water.

5. Acknowledgements

This research was part-funded by the European Social Fund (ESF) through the European Union's Convergence programme administered by the Welsh Government. Match funding and access to water samples and water quality data was provided by Dŵr Cymru Welsh Water (DCWW). Christopher Freeman and Nathalie Fenner acknowledge funding from NERC under the first EU ERA-EnvHealth call (FP7-ENV-2007- CSA-1.2.3-01) and the large NERC grant, *Characterisation of the nature, origins and ecological significance of dissolved organic matter in freshwater ecosystems* (NE/K010689/1).

6. References

- Aitkenhead, J.A., Hope, D. and Billett, M.F. (1999) 'The relationship between dissolved organic carbon in stream water and soil organic carbon pools at different spatial scales', *Hydrological Processes*, Vol. 13 No. 8, pp.1289 - 1302.
- Alfredsson, H., Condron, L.M., Clarholm, M. and Davis, M.R. (1998) 'Changes in soil acidity and organic matter following the establishment of conifers on former grassland in New Zealand', *Forest Ecology and Management*, Vol. 112 No. 3, pp. 245 - 252.
- Algesten, G., Sobek, S., Bergström, A., Ågren, A., Tranvik, L.J. and Jansson, M. (2004) 'Role of lakes for organic carbon cycling in the Boreal zone', *Global Change Biology*, Vol. 10 No. 1, pp. 141 - 147.
- Armstrong, A., Holden, J., Kay, P., Chapman, P., Clements, S., Foulger, M., McDonald, A.T. and Walker, A. (2007) *Grip-blocking in upland catchments; costs and benefits*, Tech. Rep. Final Report for Yorkshire Water.
- Badruzzaman, M., Pinzon, J., Oppenheimer, J. and Jacangelo, J.G. (2012) 'Sources of nutrients impacting surface waters in Florida: a review', *Journal of Environmental Management*, Vol. 109, pp. 80 - 92.
- Bernhardt, H., Schell, H., Hoyer, O. and Lusse, B. (1991) 'Influence of algogenic organic substances on flocculation and filtration', *WISA*, Vol. 1, pp. 41 - 57.
- Billett, M.F. and Cresser, M.S. (1992) 'Predicting stream-water quality using catchment and soil chemical characteristics', *Environmental Pollution*, Vol. 77 No. 2-3, pp. 263 - 268.
- Boddy, E., Roberts, P., Hill, P.W., Farrar, J. and Jones, D.L. (2008) 'Turnover of low molecular weight dissolved organic C (DOC) and microbial C exhibit different temperature sensitivities in Arctic tundra soils', *Soil Biology and Biochemistry*, Vol. 40 No.7, pp. 1557 - 1566.
- Campbell, C., Lafond, G., Biederbeck, V., Wen, G., Schoenau, J. and Hahn, D. (1999) 'Seasonal trends in soil biochemical attributes: effects of crop management on a black chernozem', *Canadian Journal of Soil Science*, Vol. 79 No. 1, pp. 85 - 97.
- Chantigny, M.H. (2003) 'Dissolved and water-extractable organic matter in soils: a review on the influence of land use and management practices', *Geoderma*, Vol. 113 No. 3-4, pp. 357 - 380.
- Chow, C.W., Fabris, R., Drikas, M. and Holmes, M. (2005) 'A case study of treatment performance and organic character', *Journal of Water Supply Research and Technology – AQUA*, Vol. 54 No. 6, pp. 385 - 395.
- Chow, A.T., Lee, S., O'Geen, A.T., Orozco, T., Beaudette, D., Wong, P., Hernes, P.J., Tate, K.W. and Dahlgren, R.A. (2009) 'Litter contributions to dissolved organic matter and disinfection byproduct precursors in California Oak woodland watersheds', *Journal of Environmental Quality*, Vol. 38 No. 6, pp. 2334 - 2343.

- 520 Christophersen, N. and Wright, R.F. (1981) 'Sulfate budget and a model for sulfate concentrations in
521 stream water at Birkenes, a small forested catchment in southernmost Norway', *Water Resources*
522 *Research*, Vol. 17 No. 2, pp. 377 - 389.
- 523 Clair, T.A., Pollock, T.L. and Ehrman, J.M. (1994) 'Exports of carbon and nitrogen from river basins in
524 Canada's Atlantic Provinces', *Global Biogeochemical Cycles*, Vol. 8 No. 4, pp. 441 - 450.
- 525 Clutterbuck, B. and Yallop, A.R. (2010) 'Land management as a factor controlling dissolved organic
526 carbon release from upland peat soils 2: changes in DOC productivity over four decades', *Science of*
527 *the Total Environment*, Vol. 408 No. 24, pp. 6179 - 6191.
- 528 Clymo, R. (1964) 'The origin of acidity in sphagnum bogs', *The Bryologist*, Vol. 67 No. 4, pp. 427 - 431.
- 529 Cooke, J.G. and Cooper, A.B. (1988) 'Sources and sinks of nutrients in a New Zealand hill pasture
530 catchment III. Nitrogen', *Hydrological Processes*, Vol. 2 No. 2, pp. 135 - 149.
- 531 Correll, D.L. (1988) 'The role of phosphorus in the eutrophication of receiving waters: a review',
532 *Journal of Environmental Quality*, Vol. 27 No. 2, pp. 261 - 266.
- 533 Cosby, B., Hornberger, G., Galloway, J. and Wright, R. (1985) 'Modeling the effects of acid
534 deposition: assessment of a lumped parameter model of soil water and streamwater chemistry',
535 *Water Resources Research*, Vol. 21 No. 1, pp. 51 - 63.
- 536 Cresser, M.S. and Edwards, A. (1987) *Acidification of freshwaters*, Cambridge University Press,
537 Cambridge.
- 538 Davies, J-M., Roxborough, M. and Mazumder, A. (2004) 'Origins and implications of drinking water
539 odours in lakes and reservoirs of British Columbia, Canada', *Water Research*, Vol. 38 No. 7, pp. 1900 -
540 1910.
- 541 Delprat, L., Chassin, P., Linères, M. and Jambert, C. (1997) 'Characterization of dissolved organic
542 carbon in cleared forest soils converted to maize cultivation', *European Journal of Agronomy*, Vol. 7
543 No. 1-3, pp. 201 - 210.
- 544 DEFRA (Department of Environment, Food and Rural Affairs) (2002) *Ammonia in the UK*.
- 545 Drinan, T.J., Graham, C.T., O'Halloran, J. and Harrison, S.S.C. (2013) 'The impact of catchment conifer
546 plantation forestry on the hydrochemistry of peatland lakes', *Science of the Total Environment*, Vol.
547 443, pp. 608 - 620.
- 548 DWI (Drinking Water Inspectorate). *A brief guide to drinking water safety plans*.
549 <http://www.sswm.info/content/brief-guide-water-safety-plans> (Accessed 11 August 2014).
- 550 Eckhardt, B. and Moore, T. (1990) 'Controls on dissolved organic carbon concentrations in streams,
551 southern Quebec', *Canadian Journal of Fisheries and Aquatic Sciences*, Vol. 47 No. 8, pp. 1537 - 1544.
- 552 EDINA. *Digimap collections*. [online] <http://edina.ac.uk/digimap> (Accessed 11 August 2014).

553 Edzwald, J. (1993) 'Coagulation in drinking water treatment: particles, organics and coagulants',
554 *Water Science and Technology*, Vol. 27 No. 11, pp. 21 - 35.

555 Eikebrokk, B., Vogt, R. and Liltved, H. (2004) 'NOM increase in northern European source waters:
556 discussion of possible causes and impacts on coagulation/contact filtration processes', *Water Supply*,
557 Vol. 4 No. 4, pp. 47 - 54.

558 Eisalou, H.K., Şengönül, K., Gökbülak, F., Serengil, Y. and Uygur, B. (2013) 'Effects of forest canopy
559 cover and floor on chemical quality of water in broad leaved and coniferous forests of Istanbul,
560 Turkey', *Forest Ecology and Management*, Vol. 289, pp. 371 - 377.

561 Engstrom, D.R. (1987) 'Influence of vegetation and hydrology on the humus budgets of Labrador
562 lakes', *Canadian Journal of Fisheries and Aquatic Sciences*, Vol. 44 No. 7, pp. 1306 - 1314.

563 Evans, C.D., Monteith, D.T. and Cooper, D.M. (2005) 'Long-term increases in surface water dissolved
564 organic carbon: observations, possible causes and environmental impacts', *Environmental Pollution*,
565 Vol. 137 No. 1, pp. 55 - 71.

566 Fenner, N., Freeman, C. and Worrall, F. (2009) 'Hydrological controls on dissolved organic carbon
567 production and release from UK peatlands' in Baird, A.J. et al (Eds.), *Carbon cycling in northern*
568 *peatlands*, American Geophysical Union, Washington DC, pp. 237-249.

569 Foster, J.A. and McDonald, A.T. (2000) 'Assessing pollution risks to water supply intakes using
570 geographical information systems (GIS)', *Environmental Modelling & Software*, Vol. 15 No. 3, pp. 225
571 - 234.

572 Freeman, C., Ostle, N. and Kang, H. (2001) 'An enzymic 'latch' on a global carbon store - a shortage of
573 oxygen locks up carbon in peatlands by restraining a single enzyme', *Nature*, Vol. 409 No. 6817, pp.
574 149 - 149.

575 Freeman, A.M., Lamon, III E.C. and Stow, C.A. (2009) 'Nutrient criteria for lakes, ponds, and
576 reservoirs: a Bayesian TREED model approach', *Ecological Modelling*, Vol. 220 No. 5, pp. 630 - 639.

577 Freeman, C., Evans, C.D., Monteith, D.T., Reynolds, B. and Fenner, N. (2001) 'Export of organic
578 carbon from peat soils', *Nature*, Vol. 412 No. 6849, pp. 785 - 785.

579 Fröberg, M., Hansson, K., Kleja, D.B. and Alavi, G. (2011) 'Dissolved organic carbon and nitrogen
580 leaching from scots pine, norway spruce and silver birch stands in southern Sweden', *Forest Ecology*
581 *and Management*, Vol. 262 No. 9, pp. 1742 - 1747.

582 Gao, J.P., Maguhn, J., Spitzauer, P. and Kettrup, A. (1998) 'Sorption of pesticides in the sediment of
583 the Teufelsweiher pond (southern Germany). II: competitive adsorption, desorption of aged residues
584 and effect of dissolved organic carbon', *Water Research*, Vol. 32 No. 7, pp. 2089 - 2094.

585 Garrison, P.J., Greb, S.R., Knauer, D.R., Wentz, D.A., Krohelski, J.T., Bockheim, J.G., Gherini, S.A. and
586 Chen, C.W. (1987) 'Application of the ILWAS model to the northern Great Lakes states', *Lake and*
587 *Reservoir Management*, Vol. 3 No. 1, pp. 356 - 364.

- 588 Gergel, S.E., Turner, M.G. and Kratz, T.K. (1999) 'Dissolved organic carbon as an indicator of the scale
589 of watershed influence on lakes and rivers', *Ecological Applications*, Vol. 9 No. 4, pp. 1377 - 1390.
- 590 Gough, R., Holliman, P.J., Willis, N., Jones, T.G. and Freeman, C. (2012) 'Influence of habitat on the
591 quantity and composition of leachable carbon in the O2 horizon: potential implications for potable
592 water treatment', *Lake and Reservoir Management*, Vol. 28 No. 4, pp. 282 - 292.
- 593 Grayson, R. and Western, A. (2001) 'Terrain and the distribution of soil moisture', *Hydrological
594 Processes*, Vol. 15 No. 13, pp. 2689 - 2690.
- 595 Grayson, R., Kay, P., Foulger, M. and Gledhill, S. (2012) 'A GIS based MCE model for identifying water
596 colour generation potential in UK upland drinking water supply catchments', *Journal of Hydrology*,
597 Vol. 420-421, pp. 37 - 45.
- 598 Gregorich, E., Rochette, P., McGuire, S., Liang, B. and Lessard, R. (1998) 'Soluble organic carbon and
599 carbon dioxide fluxes in maize fields receiving spring-applied manure', *Journal of Environmental
600 Quality*, Vol. 27 No. 1, pp. 209 - 214.
- 601 Hecky, R. and Kilham, P. (1988) 'Nutrient limitation of phytoplankton in freshwater and marine
602 environments: a review of recent evidence on the effects of enrichment', *Limnology and
603 Oceanography*, Vol. 33 No. 4, pp. 796 - 822.
- 604 Hejzlar, J., Dubrovský, M., Buchtele, J. and Ruzicka, M. (2003) 'The apparent and potential effects of
605 climate change on the inferred concentration of dissolved organic matter in a temperate stream (the
606 Malše river, south Bohemia)', *Science of the Total Environment*, Vol. 310 No. 1-3, pp. 143 - 152.
- 607 Hill, A. (1979) 'Denitrification in the nitrogen budget of a river ecosystem', *Nature*, Vol. 281 No.
608 5729, pp. 291 - 292.
- 609 Hindar, A., Wright, R.F., Nilsen, P., Larssen, T. and Høgberget, R. (2003) 'Effects on stream water
610 chemistry and forest vitality after whole-catchment application of dolomite to a forest ecosystem in
611 southern Norway', *Forest Ecology and Management*, Vol. 180 No. 1-3, pp. 509 - 525.
- 612 Holden, J., Chapman, P.J., Labadz, J.C. (2004) 'Artificial drainage of peatlands: hydrological and
613 hydrochemical process and wetland restoration', *Progress in Physical Geography*, Vol. 28 No. 1, pp.
614 95 - 123.
- 615 Holden, J., Shotbolt, L., Bonn, A., Burt, T.P., Chapman, P.J., Dougill, A.J., Fraser, E.D.G., Hubacek, K.,
616 Irvine, B., Kirkby, M.J., Reed, M.S., Prell, C., Stagl, S., Stringer, L.C., Turner, A. and Worrall, F. (2007)
617 'Environmental change in moorland landscapes', *Earth-Science Reviews*, Vol. 82 No. 1-2, pp. 75 - 100.
- 618 Hongve, D. (1999) 'Production of dissolved organic carbon in forested catchments', *Journal of
619 Hydrology*, Vol. 224, pp. 91 - 99.
- 620 Hope, D., Billett, M.F. and Cresser, M.S. (1994) 'A review of the export of carbon in river water:
621 fluxes and processes', *Environmental Pollution*, Vol. 84 No. 3, pp. 301 - 324.

- 622 Hope, D., Billett, M.F. and Cresser, M.S. (1997) 'Exports of organic carbon in two river systems in NE
623 Scotland', *Journal of Hydrology*, Vol. 193 No. 1-4, pp. 61-82.
- 624 Hopkins, D., Ibrahim, D., O'donnell, A. and Shiel, R. (1990) 'Decomposition of cellulose, soil organic
625 matter and plant litter in a temperate grassland soil', *Plant and Soil*, Vol. 124 No. 1, pp. 79 - 85.
- 626 Horton, R.E. (1933) 'The role of infiltration in the hydrologic cycle', *Transactions American
627 Geophysical Union*, Vol. 14, pp. 446 - 460.
- 628 Howe, L., Blackstock, T., Burrows, C. and Stevens, J. (2005) 'The habitat survey of Wales', *British
629 Wildlife*, Vol. 16 No. 3, pp. 153 – 162.
- 630 Hurley, J.P., Benoit, J.M., Babiarz, C.L., Shafer, M.M., Andren, A.W., Sullivan, J.R., Hammond, R. and
631 Webb, D.A. (1995) 'Influences of Watershed Characteristics on Mercury Levels in Wisconsin Rivers',
632 *Environmental Science and Technology*, Vol. 29 No. 7, pp. 1867 - 1875.
- 633 Kalbitz, K., Solinger, S., Park, J.H., Michalzik, B. and Matzner, E. (2000) 'Controls on the dynamics of
634 dissolved organic matter in soils: a review', *Soil Science*, Vol. 165 No. 4, pp. 277 - 304.
- 635 Kawasaki, M., Ohte, N. and Katsuyama, M. (2005) 'Biogeochemical and hydrological controls on
636 carbon export from a forested catchment in central Japan', *Ecological Research*, Vol. 20 No. 3, pp.
637 347 - 358.
- 638 Lake, I.R., Lovett, A.A., Hiscock, K.M., Betson, M., Foley, A., Sünnerberg, G., Evers, S. and Fletcher, S.
639 (2003) 'Evaluating factors influencing groundwater vulnerability to nitrate pollution: developing the
640 potential of GIS', *Journal of Environmental Management*, Vol. 68 No. 3, pp. 315 - 328.
- 641 Levine, D.A. and Jones, W.W. (1990) 'Modeling phosphorus loading to three Indiana reservoirs: a
642 geographic information system approach', *Lake and Reservoir Management*, Vol. 6 No. 1, pp. 81 - 91.
- 643 Li, L., Gao, N., Deng, Y., Yao, J., Zhang, K. (2012) 'Characterization of intracellular & extracellular
644 algae organic matters (AOM) of microcystic aeruginosa and formation of AOM-associated
645 disinfection byproducts and odor & taste compounds', *Water Research*, Vol. 46 No. 4, pp. 1233 -
646 1240.
- 647 Lumsdon, D.G., Stutter, M.I., Cooper, R.J. and Manson, J.R. (2005) 'Model assessment of
648 biogeochemical controls on dissolved organic carbon partitioning in an acid organic soil',
649 *Environmental Science and Technology*, Vol. 39 No. 20, pp. 8057 - 8063.
- 650 Marrs, R.H., Le Duc, M.G., Mitchell, R.J., Goddard, D., Paterson, S. and Pakeman, R.J. (2000) 'The
651 ecology of bracken: its role in succession and implications for control', *Annals of Botany*, Vol. 85
652 (suppl B), pp. 3 - 15.
- 653 Martin, C., Aquilina, L., Gascuel-Oudou, C., Molénat, J., Faucheux, M. and Ruiz, L. (2004) 'Seasonal
654 and interannual variations of nitrate and chloride in stream waters related to spatial and temporal

655 patterns of groundwater concentrations in agricultural catchments', *Hydrological Processes*, Vol. 18
656 No. 7, pp. 1237 - 1254.

657 McFee, W.W. and Kelly, J.M. (1995) 'Management-induced changes in the actively cycling fractions
658 of soil organic matter', *Carbon*, Vol. 1, pp. 119 - 138.

659 Miller, C.J. (2008) *Mechanisms of water colour release from organic soils and consequences for*
660 *catchment management*. Unpublished PhD thesis, University of Aberdeen, Aberdeen.

661 MAFF (Ministry of Agriculture, Fisheries and Food) (2000) *Fertiliser recommendations for agricultural*
662 *and horticultural crops (RB209)*.

663 Mitchell, G. and McDonald, A. (1992) 'Relationship between different methods of colour
664 measurement in potable water', *Transactions of the Institute of Measurement and Control*, pp. 306 -
665 309.

666 Mitchell, G.N. (1991) 'Water quality issues in the British uplands', *Applied Geography*, Vol. 11 No. 3,
667 pp. 201 - 214.

668 Mitsch, W. and Gosselink, J. (2007) *Wetlands*, 4th ed., John Wiley & Sons, New Jersey.

669 Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, H.A., Forsius, M., Hogasen, T., Wilander, A.,
670 Skjelkvale, B.L., Jeffries, D.S., Vuorenmaa, J., Keller, B., Kopacek, J. and Vesely, J. (2007) 'Dissolved
671 organic carbon trends resulting from changes in atmospheric deposition chemistry', *Nature*, Vol.
672 450, pp. 537 - 540.

673 Neill, M. (1989) 'Nitrate concentrations in river waters in the south-east of Ireland and their
674 relationship with agricultural practice', *Water Research*, Vol. 23 No. 11, pp. 1339 - 1355.

675 Nguyen, M.L., Westerhoff, P., Baker, L., Hu, Q., Esparza-Soto, M. and Sommerfeld, M. (2005)
676 'Characteristics and reactivity of algae-produced dissolved organic carbon', *Journal of Environmental*
677 *Engineering*, Vol. 131 No. 11, pp. 1574 - 1582.

678 Nykvist, N. (1963) 'Leaching and decomposition of water-soluble organic substances from different
679 types of leaf and needle litter', *Studia Forestalia Suecica*, Vol. 3, pp. 1 - 31.

680 Ordóñez, J.A.B., de Jong, B.H.J., García-Oliva, F., Aviña, F.L., Pérez, J.V., Guerrero, G., Martínez, R. and
681 Masera, O. (2008) 'Carbon content in vegetation, litter, and soil under 10 different land-use and
682 land-cover classes in the central highlands of Michoacan, Mexico', *Forest Ecology and Management*,
683 Vol. 255 No. 7, pp. 2074 - 2084.

684 Osborne, L.L. and Wiley, M.J. (1988) 'Empirical relationships between land use/cover and stream
685 water quality in an agricultural watershed', *Journal of Environmental Management*, Vol. 26 No. 1, pp.
686 9 - 27.

- 687 Potthast, K., Hamer, U. and Makeschin, F. (2012) 'Land-use change in a tropical mountain rainforest
688 region of southern Ecuador affects soil microorganisms and nutrient cycling', *Biogeochemistry*, Vol.
689 111 No. 1-3, pp. 151 - 167.
- 690 Prévost, M., Rompré, A., Coallier, J., Servais, P., Laurent, P., Clément, B. and Lafrance, P. (1998)
691 'Suspended bacterial biomass and activity in full-scale drinking water distribution systems: impact of
692 water treatment', *Water Research*, Vol. 32 No. 5, pp. 1393 - 1406.
- 693 Rasmussen, L.H., Kroghsbo, S., Frisvad, J.C. and Hansen, H.C.B. (2003) 'Occurrence of the
694 carcinogenic bracken constituent ptaquiloside in fronds, topsoils and organic soil layers in Denmark',
695 *Chemosphere*, Vol. 51 No. 2, pp. 117 - 127.
- 696 Rasmussen, J.B., Godbout, L. and Schallenberg, M. (1989) 'The humic content of lake water and its
697 relationship to watershed and lake morphometry', *Limnology and Oceanography*, Vol. 34 No. 7, pp.
698 1336 - 1343.
- 699 Reynolds, B., Emmett, B.A. and Woods, C. (1992) 'Variations in streamwater nitrate concentrations
700 and nitrogen budgets over 10 years in a headwater catchment in mid-Wales', *Journal of Hydrology*,
701 Vol. 136 No. 1-4, pp. 155 - 175.
- 702 Ritson, J.P., Graham, N.J.D., Templeton, M.R., Clark, J.M., Gough, R. and Freeman, C. (2014) 'The
703 impact of climate change on the treatability of dissolved organic matter (DOM) in upland water
704 supplies: a UK perspective', *Science of the Total Environment*, Vol. 473-474 No. 1, pp. 714 - 730.
- 705 Rochelle, B.P., Liff, C.I., Campbell, W.G., Cassell, D.L., Robbins Church, M. and Nusz, R.A. (1989)
706 'Regional relationships between geomorphic/hydrologic parameters and surface water chemistry
707 relative to acidic deposition', *Journal of Hydrology*, Vol. 112 No. 1, pp. 103 - 120.
- 708 Rothwell, J.J., Evans, M.G., Daniels, S.M. and Allott, T.E.H. (2007) 'Baseflow and stormflow metal
709 concentrations in streams draining contaminated peat moorlands in the Peak District National Park
710 (UK)', *Journal of Hydrology*, Vol. 341 No. 1-2, pp. 90 - 104.
- 711 Scott, M.J., Jones, M.N., Woof, C. and Tipping, E. (1998) 'Concentrations and fluxes of dissolved
712 organic carbon in drainage water from an upland peat system', *Environment International*, Vol. 24
713 No. 5-6, pp. 537 - 546.
- 714 Skjelkvåle, B.L., Stoddard, J.L., Jeffries, D.S., Tørseth, K., Høgåsen, T., Bowman, J., Mannio, J.,
715 Monteith, D.T., Mosello, R., Rogora, M., Rzychon, D., Vesely, J., Wieting, J., Wilander, A. and
716 Worsztynowicz, A. (2005) 'Regional scale evidence for improvements in surface water chemistry
717 1990-2001', *Environmental Pollution*, Vol. 137 No. 1, pp. 165 - 176.
- 718 Smith, V.H. (1998) 'Cultural eutrophication of inland, estuarine, and coastal waters' in Pace, M.L. et
719 al (Eds.), *Successes, Limitations, and Frontiers in Ecosystem Science*, Springer-Verlag, New York, pp. 7
720 - 49.

- 721 Sobek, S., Tranvik, L.J., Prairie, Y.T., Kortelainen, P. and Cole, J.J. (2007) 'Patterns and regulation of
722 dissolved organic carbon: an analysis of 7,500 widely distributed lakes', *Limnology and*
723 *Oceanography*, Vol. 52 No. 3, pp. 1208 - 1219.
- 724 Soulsby, C., Tetzlaff, D., Rodgers, P., Dunn, S. and Waldron, S. (2006) 'Runoff processes, stream water
725 residence times and controlling landscape characteristics in a mesoscale catchment: an initial
726 evaluation', *Journal of Hydrology*, Vol. 325 No. 1-4, pp. 197 - 221.
- 727 Stevens, P.A., Hornung, M. and Hughes, S. (1989) 'Solute concentrations, fluxes and major nutrient
728 cycles in a mature sitka-spruce plantation in Beddgelert forest, north Wales', *Forest Ecology and*
729 *Management*, Vol. 27 No. 1, pp. 1 - 20.
- 730 Stevens, J.P., Blackstock, T.H., Howe, E.A. and Stevens, D.P. (2004) 'Repeatability of Phase 1 habitat
731 survey', *Journal of Environmental Management*, Vol. 73 No. 1, pp. 53-59.
- 732 Stoddard, J.L., Kahl, J.S., Deviney, F.A., Dewalle, D.R., Driscoll, C.T. and Herlihy, A.T. (2003) *Response*
733 *of surface water chemistry to the clean air act amendments of 1990*, Tech. Rep. EPA 620/R-03/001.
- 734 Stutter, M.I., Deeks, L.K., Low, D. and Billett, M.F. (2006) 'Impact of soil and groundwater
735 heterogeneity on surface water chemistry in an upland catchment', *Journal of Hydrology*, Vol. 318
736 No. 1-4, pp. 103 - 120.
- 737 Thurman, E.M. (Ed.), (1985) *Organic Geochemistry of Natural Waters*, Kluwer Academic Publishers,
738 Lancaster.
- 739 Tipping, E., Woof, C., Rigg, E., Harrison, A.F., Ineson, P., Taylor, K., Benham, D., Poskitt, J., Rowland,
740 A.P., Bol, R. and Harkness, D.D. (1999) 'Climatic influences on the leaching of dissolved organic
741 matter from upland UK moorland soils, investigated by a field manipulation experiment',
742 *Environment International*, Vol. 25 No. 1, pp. 83 - 95.
- 743 Toberman, H., Evans, C.D., Freeman, C., Fenner, N., White, M., Emmett, B.A. and Artz, R.R.E. (2008)
744 'Summer drought effects upon soil and litter extracellular phenol oxidase activity and soluble carbon
745 release in an upland *calluna* heathland', *Soil Biology and Biochemistry*, Vol. 40 No. 6, pp. 1519 -
746 1532.
- 747 Trudgill, S., Burt, T., Heathwaite, A.L. and Arkell, B. (1991) 'Soil nitrate sources and nitrate leaching
748 losses, Slapton, south Devon', *Soil Use and Management*, Vol. 7 No. 4, pp. 200 - 206.
- 749 Urban, N.R., Verry, E.S. and Eisenreich, S.J. (1995) 'Retention and mobility of cations in a small
750 peatland - trends and mechanisms', *Water, Air, and Soil Pollution*, Vol. 79 No. 1-4, pp. 201 - 224.
- 751 van den Berg, L.J.L., Shotbolt, L. and Ashmore, M.R. (2012) 'Dissolved organic carbon (DOC)
752 concentrations in UK soils and the influence of soil, vegetation type and seasonality', *Science of the*
753 *Total Environment*, Vol. 427-428, pp. 269 - 276.

754 Vollenweider, R.A. (1968) *The scientific fundamentals of lake and stream eutrophication, with*
755 *particular reference to phosphorus and nitrogen as eutrophication factors*, Tech. Rep. DAS/
756 DSI/68.27.

757 Wallage, Z.E., Holden, J. and McDonald, A.T. (2006) 'Drain blocking: an effective treatment for
758 reducing dissolved organic carbon loss and water discolouration in a drained peatland', *Science of*
759 *the Total Environment*, Vol. 367 No. 2-3, pp. 811 - 821.

760 Watts, C.D., Naden, P.S., Machell, J. and Banks, J. (2001) 'Long term variation in water colour from
761 Yorkshire catchments', *Science of the Total Environment*, Vol. 278 No. 1-3, pp. 57 - 72.

762 Weiler, M. and Naef, F. (2003) 'Simulating surface and subsurface initiation of macropore flow',
763 *Journal of Hydrology*, Vol. 273 No. 1, pp. 139 - 154.

764 WHO (World Health Organization). (2011) *Guidelines for drinking-water quality*.

765 Worrall, F., Reed, M., Warburton, J. and Burt, T. (2003) 'Carbon budget for a British upland peat
766 catchment', *Science of the Total Environment*, Vol. 312 No. 1-3, pp. 133 - 146.

767 Zhang, W., An, S., Xu, Z., Cui, J. and Xu, Q. (2011) 'The impact of vegetation and soil on runoff
768 regulation in headwater streams on the east Qinghai-Tibet Plateau, China', *CATENA*, Vol. 87 No. 2,
769 pp. 182 - 189.

770 Zsolnay, A. (1996) 'Dissolved humus in soil waters', in Piccolo, A. (Ed.), *Humic Substances in*
771 *Terrestrial Ecosystems*, Elsevier Science, Amsterdam, pp. 171 - 223.

772

7. Tables and Figures

Table 1. Categorisation of Phase 1 habitat types.

Category used in present study	Phase 1 Habitat classification	Category used in present study	Phase 1 Habitat classification
Woodland and scrub	A.1 A.2	Fen/ mire	E.3
Recently-felled woodland	A.4	Bare peat	E.4
Unimproved grassland	B.1.1 B.3.1	Swamp	F.1 F.2
Improved grassland	B.1.2 B.2.2 B.4	Water	G.1 G.2
Marsh/ marshy grassland	B.5	Rock/ scree/ quarry	I.1 I.2
Tall herb and fern	C.1 C.2 C.3	Arable	J.1.1
Heathland	D.1 D.2 D.3 D.5 D.6	Caravan site	J.3.4
Bog	E.1	Buildings	J.3.6
Flush and spring	E.2	Other	J.1.2 J.4 Not accessed Illegible

Table 2. Correlation coefficients (r) for statistically significant correlations between percentage Phase 1 Habitat coverage and reservoir water quality (whole-catchment analysis).

	Autumn [DOC]	Spring [DOC]	Autumn Colour (Hazen)	Spring Colour (Hazen)	Autumn [NO ₃]	Spring [NO ₃]	Autumn pH	Spring pH
Woodland and scrub						0.743**		
Recently-felled woodland								
Improved grassland					0.636**	0.677**		
Unimproved grassland		-0.512*						
Marsh/ marshy grassland								
Tall herb and fern	-0.499*		-0.588*					
Heathland	-0.543*		-0.543*	-0.649**				
Bog								
Flush and spring								
Fen/ mire	0.564*					0.577*		
Bare peat								
Swamp	0.612*	0.690**	0.624**	0.636**				
Water								
Rock/ scree/ quarry								
Arable	0.734**	0.651**	0.508*	0.549*	0.632**	0.721**		
Caravan site								
Buildings	0.596*					0.580*		
Other	0.548*			0.539*				

* indicates $p < 0.05$ and ** indicates $p < 0.01$. All results shown relate to Spearman's correlation analysis.

Table 3. Correlation coefficients (*r*) for statistically significant correlations between percentage Phase 1 Habitat coverage and reservoir water quality (250 m buffer zone analysis).

	Autumn [DOC]	Spring [DOC]	Autumn Colour (Hazen)	Spring Colour (Hazen)	Autumn [NO ₃]	Spring [NO ₃]	Autumn pH	Spring pH
Woodland and scrub						0.774**		
Recently-felled woodland								
Improved grassland					0.617*	0.655**		
Unimproved grassland								
Marsh/ marshy grassland						0.502*		
Tall herb and fern			-0.559*					
Heathland	-0.560*	<u>-0.517*</u>	-0.499*	-0.652**			-0.558*	
Bog								
Flush and spring								
Fen/ mire	0.600*	0.513*			0.587*	0.578*		
Bare peat								
Swamp	0.647**	0.709**	0.624**	0.636**				
Water								
Rock/ scree/ quarry								
Arable								
Buildings	0.499*							
Other								

* indicates $p < 0.05$ and ** indicates $p < 0.01$. Underlined result relates to Pearson's correlation analysis and the remainder to Spearman's correlation analysis.




Figure 1. Percentage Phase 1 Habitat coverage in 250 m reservoir buffer zone.




Figure 2. Percentage Phase 1 Habitat coverage in whole reservoir catchments.